

## DEFINITION

Otitis Externa (OE) is an infection or diffuse inflammation of the ear canal lining which presents in two forms:

1. A benign painful infection of the outer canal.
2. Malignant (necrotizing) OE: A potentially lethal form that usually occurs in immunocompromised or diabetic clients. It involves bacterial spread to the cartilage of the external ear with pain and edema. It may be accompanied by a fever and systemic manifestations of infection.

OE is also called swimmer's ear, tropical ear, and external ear infection.

## IMMEDIATE CONSULTATION REQUIRED IN THE FOLLOWING SITUATIONS

- Fever > 39°C oral
- Immunocompromised
- Extensive cervical lymphadenopathy
- Mastoid process erythema, tenderness, swelling with fever, protrusion of pinna
- Facial nerve palsy
- Suspicions of malignant OE
- Initial therapy failure or recurrence
- Local cellulitis
- Parotitis
- Vertigo

Types:

Acute

- Usually due to bacterial infection (swimmer's ear)
- Most common
- Sterile canal or Staphylococcus found in 70-80% of children
- Severe infection could be due to Pseudomonas infection

Chronic

- Lasting > 3 months
- Usually fungal or allergic in origin
- May be manifestation of dermatitis
- May result from incomplete resolution of acute form

OTITIS EXTERNA PEDIATRIC

Localized

- Due to folliculitis or furuncle in external ear canal
- Occurs usually in outer third of canal

Diffuse OE

- Due to more widespread inflammation of the skin and subdermis of the external ear canal

Malignant OE

- Also called malignant external otitis, necrotizing external otitis, necrotizing OE, or osteomyelitis of ear canal.
- Extends beyond external auditory canal and may include the mastoid bone, cranial nerves (facial nerve with glossopharyngeal and less often spinal accessory nerves), parotid gland, paranasal sinuses, veins, or meninges. *Pseudomonas aeruginosa* (*P. aeruginosa*) is usually the organism involved.

**CAUSES**

A disruption of the skin or cerumen protective barrier in the external ear canal leads to OE.

Acute OE

- Gram-negative rods: Proteus, Pseudomonas
- Gram-positive cocci (less common): Staphylococcus, Streptococcus
- Fungal infection (e.g., Aspergillus and Candida)

Chronic OE

- Inadequate treatment of acute form
- Allergic contact dermatitis
- Seborrheic dermatitis
- Food sensitivity associated with atopic dermatitis
- Systemic dermatitis
- Generalized skin conditions (atopic dermatitis/eczema, psoriasis)
- Variants of chronic OE
- Type IV cell-mediated hypersensitivity reaction to ototopical treatment of acute form
- Tympanostomy tube leading to middle ear drainage causing canal maceration

**OTITIS EXTERNA PEDIATRIC**

- Dermatophytid eruptions caused by hematogenous spread of fungal by-products from primary site external to ear

Localized OE

- Typically caused by infection of hair follicle by *Staphylococcus aureus* (*S. aureus*)

Malignant OE

- *P. aeruginosa*
- *S. aureus*
- Proteus species
- Klebsiella

**PREDISPOSING AND RISK FACTORS**

- Hearing aids
- Narrow ear canal
- Use of cotton-tipped applicators
- Use of ear plugs
- Swimming
- Excessive bathing
- Histocytosis
- Immunocompromised conditions

Likely risk factors:

Activities/conditions that introduce irritation, trauma, or infectious agent:

- Prolonged swimming
- Hearing aids
- Impacted cerumen
- Insufficient cerumen
- Microtrauma secondary to ear cleaning

Conditions that disrupt epithelium of the skin canal such as:

- Seborrheic dermatitis
- Eczema
- Psoriasis

**OTITIS EXTERNA PEDIATRIC**

Possible risk factors:

- High humidity
- Warmer temperatures
- Exposure to water high in bacterial counts
- Chronic dermatologic conditions
- Maceration of skin
- Sweating
- Allergy
- Stress
- Anatomic abnormalities (narrow canal or exostoses) may predispose client to acute OE
- Previous ear surgery (such as tympanostomy)

Risk factors for malignant OE:

- Diabetes
- Immunocompromised state
- Radiation therapy to head and neck

**HISTORY**

- Ear pain (otalgia)
- Pruritus or irritation
- Purulent discharge from canal (cheesy white, greenish blue or grey)
- Recent exposure to water or mechanical trauma
- Reduced hearing or feelings of fullness in ear may be present
- Unilateral headache may be present
- Erythema
- Edema
- Tinnitus
- Jaw pain
- Facial paralysis, vertigo, meningeal signs (such as headache) may be present in malignant external otitis
- 90% of cases are unilateral

## PHYSICAL FINDINGS

- Temperature may be  $> 39^{\circ}\text{C}$  oral with malignant OE but not common with acute otitis externa.
- Pain with pinna/tragus manipulation or pressure may be used to differentiate OE from acute otitis media with otorrhea.
- Regional cellulitis of auricle, pinna, and adjacent skin may be present in some clients.
- Necrosis or granulation of canal skin in malignant OE.
- Look for other dermatologic manifestations (such as eczema, psoriasis).

Acute forms may have:

- Ear canal edema, erythema, or both
- Erythemic tympanic membrane
- Canal obstruction
- Aural discharge may occur, may be seropurulent
- Auricular cellulitis
- Parotitis
- Enlarged cervical lymph nodes

Chronic forms may have:

- Erythema of external canal
- Lichenification of skin of external canal
- Whitish cotton-like strands in canal (with or without black or white fungal balls) with fungal infections
- Canal appearance with “wet newspaper” with grey or slightly black colour
- Check for tympanic membrane integrity
- Ear manipulation may be painful, especially pressure on tragus or movement of auricle
- Cervical adenopathy may be present with severe disease and extra canal manifestations

## DIFFERENTIAL DIAGNOSIS

- Acute otitis media with perforation

**OTITIS EXTERNA PEDIATRIC**

- Chronic suppurative otitis media
- Skin condition involving the ear (e.g., eczema)
- Mastoiditis
- Furuncle in canal
- Foreign body irritation
- Ear damage due to radiation treatment
- Acute mastoiditis
- Carcinoma of the external auditory canal
- Ramsay Hunt syndrome
- Apocrine hidrocystoma of external auditory canal
- Bullous myringitis

**COMPLICATIONS**

- Ear canal narrowing
- Tympanic membrane perforation
- Conductive hearing loss
- Extracanalicular dissemination of infection (auricular cellulitis, chondritis, parotitis, adenopathy)
- Opportunistic bacterial or fungal infection may complicate chronic OE
- Fungal involvement uncommon in primary OE but may be more common in chronic OE or after treatment with topical antibiotics (or less often, systemic antibiotics)
- Furuncle
- Regional lymphadenitis
- Progression to malignant OE may rarely lead to osteomyelitis of ear canal, perichondritis, bacterial meningitis)
- External auditory canal cholesteatoma
- Mastoiditis

**INVESTIGATIONS AND DIAGNOSTIC TESTS**

Otoscopy

- Will reveal diffuse ear canal edema, erythema, or both, with or without otorrhea or material in ear canal

**OTITIS EXTERNA PEDIATRIC**

Pneumatic otoscopy findings

- Good tympanic membrane mobility with acute OE
- Absent or limited mobility with acute otitis media and associated middle-ear effusion

Tympanometry findings

- Normal peaked curve with acute OE
- Flat tracing with acute otitis media

Culture of ear canal can identify fungi, resistant bacteria, or unusual causes of infection. Testing is usually unnecessary.

**MAKING THE DIAGNOSIS**

Criteria for diagnosis of acute OE:

- Rapid onset (usually within 48 hours) in past 3 weeks
- Symptoms of ear canal inflammation:
  - Ootalgia (often severe)
  - Itching
  - Fullness
  - May or may not have hearing loss or jaw pain
- Signs of ear canal inflammation:
  - Tenderness of tragus and/or pinna
  - Diffuse ear canal edema and/or erythema
  - May or may not have otorrhea, regional lymphadenitis, tympanic membrane erythema, or cellulitis of pinna and adjacent skin
- Consider malignant OE if:
  - Presence of severe external otitis
  - Intractable otalgia and/or headache
  - Pain and tenderness near ear and mastoid
  - Persistent discharge from ear
  - Granulation tissue at junction of bone and cartilage in external ear

Rule out:

- Unusual causes of OE

**OTITIS EXTERNA PEDIATRIC**

- Viral infections of external ear
- Otomycosis (fungal infection of external ear)
- Contact dermatitis of ear canal (may be cause of chronic OE)
- Secondary contact otitis (sensitization to topical treatment) (may be cause of chronic OE)
- Otitis media
  - Acute otitis media
  - Chronic suppurative otitis media
  - Otitis media in clients with perforation or tympanostomy tubes
- Foreign body in external ear
- Furunculosis
- Ear damage due to radiation therapy (may involve pinna, external canal, and periauricular region)
- Acute mastoiditis
- Carcinoma of external auditory canal
- Ramsay Hunt syndrome
- Malignant OE
- Apocrine hidrocystoma of external auditory canal

Testing overview:

- Testing is usually unnecessary
- Procedures which may distinguish OE from otitis media
  - Otoscopy, including pneumatic otoscopy
  - Tympanometry (but may not be possible due to client discomfort)

**MANAGEMENT AND INTERVENTIONS**

**Goals of Treatment**

- Relieve pain
- Prevent recurrence
- Eradicate infection

### **Appropriate Consultation**

- Usually not necessary if condition is uncomplicated.

### **Non-Pharmacological Interventions**

- Debriding the canal is critical. The importance of this step cannot be overemphasized.
- Clean the outer ear and the canal with normal saline and gently debride the area of debris and exudate with a gauze wick.
- If there is significant drainage or if there is threat of further narrowing, an ear wick (1 inch [2.5 cm] of cotton or gauze) threaded gently into the canal and left there will help keep the canal open and ensure that medicated drops reach the distal part of the canal.
- The wick will eventually fall out as edema subsides or can be removed after 2-3 days.

### **Pharmacological Interventions**

- A combination of antibiotic drops with 3% acetic acid and steroid is more effective than using individual agents alone for treating OE.
- Ear wicks may be necessary in swollen canals for the penetration of medicine into the canal. If an ear wick is used:
  - Remove after 48 hours and inspect canal
  - Replace wick if canal is still significantly narrowed
  - Wick is no longer necessary when canal has returned to almost normal size

### **Duration of Treatment:**

- Analgesia for several days may be necessary
- Antibiotic/antiseptic treatment for 7-10 days

### **Manage pain with simple analgesics:**

- Acetaminophen (Tylenol) 15 mg/kg/dose orally q4-6h prn (maximum dose 75 mg/kg/day)
- Or
- Ibuprofen 10 mg/kg/dose orally q6h prn (maximum dose 40 mg/kg/day)

OTITIS EXTERNA PEDIATRIC

As OE can be very painful, stronger analgesia may be necessary if acetaminophen or ibuprofen does not control pain.

OE (Acute Uncomplicated):

- If there is no danger of perforated tympanic membrane, start:
  - Gramicidin/polymyxin (Optimycin) eye/ear solution 4 drops qid for 7 days
- If the tympanic membrane cannot be visualized or is perforated:  
Children > 1 year of age:
  - Ciprofloxacin/dexamethasone (Ciprodex) otic solution 4 drops bid for 7 days
  - Or
  - Ciprofloxacin/hydrocortisone (CiproHC) 3 drops bid for 7 days
- Consult a physician/RN(NP) for children < 1 year of age

For recurrent episode, start the client on prophylactic measures:

- A solution of 3% acetic acid (mix 100 mL of white vinegar and 45 mL of water) 2-3 drops to the affected ear after swimming or showers

Malignant (Necrotizing) OE:

- Contact a physician/RN(NP) as treatment requires parenteral antibiotics with coverage for Pseudomonas species (e.g., Ciprofloxacin) in addition to hospital care.

Fungal OE (Otomycosis):

Antibiotic ear drops may worsen symptoms of fungal OE.

- Fungal organisms can cause OE especially in immunocompromised clients. The following antifungal agents can be used in mild to moderate cases of OE **but only if** the tympanic membrane is intact.
  - In mild to moderate cases of OE due to fungi, treat with antifungal agents:
    - Locacorten Vioform otic drops 2 drops bid for 7 days
- Canal should be kept free of water for at least 2 weeks.
- Canal debridement is necessary at times for resolution of infection.
- Daily meticulous irrigation with 3% acetic acid.
- Ear wicks should be used with an antifungal if tympanic membrane cannot be well visualized due to canal inflammation/debris.

**OTITIS EXTERNA PEDIATRIC**

Consult a physician/RN(NP) for severe OE that require systemic antibiotic and in those who are immunocompromised.

**Client and Caregiver Education**

- Counsel client/caregiver about appropriate use of medications (if possible, have another family member instill drops and clean the ear).
- Counsel client/caregiver about proper ear hygiene before instilling medications.
- Advise client/caregiver about preventing recurrent irritation (e.g., cotton-tipped applicators in the ears should not be used).
- Recommend proper drying of ears after swimming or use of ear plugs while swimming, bathing, or showering.
- Counsel client/caregiver about proper hygiene of hearing aids and ear plugs.
- Keep ear dry and avoid water sports for 7-10 days.
- Limit use of ear phones and hearing aids until after pain and discharge has cleared.
- Cut nails short for clients with eczema to minimize trauma.

**Monitoring and Follow-Up**

- If client fails to respond to initial therapeutic option within 48-72 hours, reassess the client to confirm diagnosis of acute OE and to rule out other causes of illness.
- Consider at least one follow-up visit to verify resolution and clear remaining debris from canal for all but mild cases.
- Instruct client to return if pain increases or if fever develops despite therapy.
- If an ear wick is used, remove after 24-48 hours and reassess.

**Referral**

- Refer to a physician/RN(NP) if treatment fails.

**DOCUMENTATION**

- As per employer policy

## REFERENCES

- Anti-Infective Review Panel. (2013). *Anti-infective guidelines for community-acquired infections*. Toronto, ON: MUMS Guideline Clearinghouse.
- Health Canada. (2011). *First Nations & Inuit health: Clinical practice guidelines for nurses in primary care*. Ottawa, ON: Author. Retrieved from <http://www.hc-sc.gc.ca>
- Hui, C. P. (2013). Acute otitis externa. *Paediatrics & Child Health (1205-7088)*, 18(2), 96–98.
- McWilliams, C. J., Smith, C. H., & Goldman, R. D. (2012). Acute otitis externa in children. *Canadian Family Physician*, 58(11), 1222–1224.
- Osguthorpe, J. D., & Shughnessy, A. F. (2013, May 30). *Otitis externa*. Retrieved from <http://www.essentialevidenceplus.com>
- Otitis externa*. (2010, May 10). Retrieved from <http://www.mdconsult.com>
- Otitis externa*. (2012, May 15). Retrieved from <http://www.pemsoft.com>
- Otitis externa*. (2013, January 07). Retrieved from <https://dynamed.ebscohost.com>
- Prentice, P. (2015). American academy of otolaryngology: Head and neck surgery foundation clinical practice guideline on acute otitis externa 2014. *Archives of Disease in Childhood - Education & Practice Edition*, 100(4), 197–197. <http://doi.org/10.1136/archdischild-2014-307676>
- What are the effects of empirical treatments for otitis externa?* (2010, August 03). Retrieved from <http://clinicalevidence.bmj.com>
- Schaefer, P., & Baugh, R. F. (2012). Acute otitis externa: An update. *American Family Physician*, 86(11), 1055–1061.

SASKATCHEWAN REGISTERED NURSES' ASSOCIATION

RNs WITH ADDITIONAL AUTHORIZED PRACTICE  
CLINICAL DECISION TOOL  
DECEMBER 1, 2016

**OTITIS EXTERNA PEDIATRIC**

NOTICE OF INTENDED USE OF THIS CLINICAL DECISION TOOL

This SRNA Clinical Decision Tool (CDT) exists solely for use in Saskatchewan by an RN with additional authorized practice as granted by the SRNA. The CDT is current as of the date of its publication and updated every three years or as needed. A member must notify the SRNA if there has been a change in best practice regarding the CDT. This CDT does not relieve the RN with additional practice qualifications from exercising sound professional RN judgment and responsibility to deliver safe, competent, ethical and culturally appropriate RN services. The RN must consult a physician/RN(NP) when clients' needs necessitate deviation from the CDT. While the SRNA has made every effort to ensure the CDT provides accurate and expert information and guidance, it is impossible to predict the circumstances in which it may be used. Accordingly, to the extent permitted by law, the SRNA shall not be held liable to any person or entity with respect to any loss or damage caused by what is contained or left out of this CDT.

SRNA © This CDT is to be reproduced only with the authorization of the SRNA.